


# Lake Okeechobee, Florida


## Sport Fish Population Evaluation

June 2005




# Electrofishing sites for largemouth bass population evaluation







**Catch rates of Largemouth bass collected via November electrofishing on Lake Okeechobee from 1992 through 2004. The blue line indicates the mean catch value and red lines depict the upper and lower 95% confidence intervals (CI).**



**Comparison of length frequency plots of largemouth bass collected with electrofishing from Lake Okeechobee. Annual sampling effort was 990 minutes (22 sites; three 15 minute repetitions at each site).**



**Comparison of length frequency plots of largemouth bass collected with electrofishing from Lake Okeechobee. Annual sampling effort was 990 minutes (22 sites; three 15 minute repetitions at each site).**




Length frequency plot of largemouth bass collected with electrofishing from Lake Okeechobee. Annual sampling effort was 990 minutes (22 sites; three 15 minute repetitions at each site)


# Trawling for black crappie

JAN 14 2003







**Catch rates of black crappie collected with 10-meter otter trawl in January from Lake Okeechobee, 1973 through 2005.**



**Catch rates of black crappie >203 mm collected with 10-meter otter trawl in January from Lake Okeechobee, 1973 through 2005.**



**Size and age structure comparison of black crappie collected with the 10-meter otter trawl from Lake Okeechobee, January 2001 to January 2004.**



**Size and age structure of black crappie collected with the 10-meter otter trawl from Lake Okeechobee, January 2005.**